



## Conversation starters for your next rheumatologist appointment

Set the appointment. Then, set yourself up for success by sharing your answers to the questions below with your rheumatologist.

Could a treatment like BIMZELX® (bimekizumab-bkzx) be right for you?  
Here are a few questions to think about:

**1** How often does your psoriatic arthritis impact your everyday life?



☐ Little to no time



☐ More than I want it to



☐ Way too much

**2** There are many symptoms of PsA— which do you regularly experience?

☐ Joint pain

☐ Joint swelling

☐ Stiffness

☐ Fatigue

☐ Swollen fingers & toes

☐ Skin plaques

**3** How does PsA regularly affect the way you feel?

☐ Increased stress

☐ Avoiding activity

☐ Skipping social activity

☐ Feeling tired

☐ Inability to sleep

☐ None apply

**4** How much does your psoriatic arthritis affect your emotional well-being?



☐ Little to no impact



☐ Medium impact



☐ Significant impact

**5** Rate how well your current treatment is working.

☐ It works well

☐ It's all right, I guess

☐ It's just not working for me

 **Bimzelx**<sup>®</sup>  
(bimekizumab-bkzx)

### USE FOR BIMZELX

BIMZELX<sup>®</sup> is a prescription medicine used to treat adults with active psoriatic arthritis.

### IMPORTANT SAFETY INFORMATION

Please see additional **Important Safety Information** on the following page and **full Prescribing Information**, including **Medication Guide**, at [BIMZELX.com](https://www.bimzelx.com).

### What is the most important information I should know about BIMZELX<sup>®</sup> (bimekizumab-bkzx)?

BIMZELX is a medicine that affects your immune system. BIMZELX may increase your risk of having serious side effects, including:

- **Suicidal thoughts and behavior have happened in some people treated with BIMZELX.** Get medical help right away or call the National Suicide and Crisis Lifeline at 988 if you, your caregiver or your family member notice in you any of the following symptoms:

## IMPORTANT SAFETY INFORMATION (cont'd)

### What is the most important information I should know about BIMZELX® (bimekizumab-bkzx)? (cont'd)

- new or worsening depression or anxiety
- thoughts of suicide, dying, or hurting yourself
- changes in behavior or mood
- acting on dangerous impulses
- attempt to commit suicide
- **Infections.** BIMZELX is a medicine that may lower the ability of your immune system to fight infections and may increase your risk of infections, including serious infections.
  - Your healthcare provider should check you for infections and tuberculosis (TB) before starting treatment with BIMZELX.
  - If your healthcare provider feels you are at risk for TB, you may be treated with medicine for TB before you begin treatment with BIMZELX and during your treatment.
  - Your doctor should watch you closely for signs and symptoms of TB during and after treatment with BIMZELX. **Do not take BIMZELX if you have an active TB infection.**

### Before starting BIMZELX, tell your healthcare provider if you:

- are being treated for an infection
- have an infection that does not go away or that keeps coming back
- have TB or have been in close contact with someone with TB
- think you have an infection or have symptoms of an infection such as:
  - fever, sweats, or chills
  - muscle aches
  - cough
  - shortness of breath
  - blood in your phlegm
  - weight loss
  - warm, red, or painful skin or sores on your body different from your psoriasis
  - diarrhea or stomach pain
  - burning when you urinate or urinating more often than normal

**After starting BIMZELX, call your healthcare provider right away if you have any of the signs of infection listed above.** Do not use BIMZELX if you have any signs of infection unless you are instructed to by your healthcare provider. See **"What are the possible side effects of BIMZELX?"** for more information about side effects.

### What is BIMZELX?

BIMZELX is a prescription medicine used to treat:

- adults with moderate-to-severe plaque psoriasis who may benefit from taking injections or pills (systemic therapy) or treatment using ultraviolet light alone or with pills (phototherapy)
- adults with active psoriatic arthritis
- adults with active non-radiographic axial spondyloarthritis with objective signs of inflammation
- adults with active ankylosing spondylitis
- adults with moderate-to-severe hidradenitis suppurativa.

It is not known if BIMZELX is safe and effective in children.

### Before using BIMZELX, tell your healthcare provider about all of your medical conditions, including if you:

- have any of the conditions or symptoms listed in the section **"What is the most important information I should know about BIMZELX?"**
- have a history of depression, or suicidal thoughts or behavior
- have liver problems
- have inflammatory bowel disease (Crohn's disease or ulcerative colitis)
- have recently received or are scheduled to receive an immunization (vaccine). You should avoid receiving live vaccines during treatment with BIMZELX.
- are pregnant or plan to become pregnant. It is not known if BIMZELX can harm your unborn baby.
  - If you become pregnant while taking BIMZELX, you are encouraged to enroll in the Pregnancy Registry, which is used to collect information about the health of you and your baby. Talk to your healthcare provider or call 1-877-311-8972 to enroll in this registry or visit <http://mothertobaby.org/pregnancy-studies/>.
- are breastfeeding or plan to breastfeed. It is not known if BIMZELX passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with BIMZELX.

**Tell your healthcare provider about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements.



**Bimzelx®**  
(bimekizumab-bkzx)

## IMPORTANT SAFETY INFORMATION (cont'd)

### What are the possible side effects of BIMZELX?

**BIMZELX may cause serious side effects. See "What is the most important information I should know about BIMZELX?"**

- **Elevated liver enzyme levels.** Your healthcare provider will do blood tests to check your liver enzyme levels before starting treatment and during treatment with BIMZELX. Your healthcare provider may temporarily stop or permanently stop your treatment with BIMZELX if you develop liver problems. Call your healthcare provider right away if you develop any signs or symptoms of liver problems, including:
  - pain on the right side of your stomach-area
  - feeling very tired
  - loss of appetite
  - nausea and vomiting
  - itching
  - dark urine
  - light-colored stool
  - yellowing of your skin or the whites of your eyes
- **Inflammatory bowel disease.** New cases of inflammatory bowel disease or "flare-ups" have happened with BIMZELX. If you have inflammatory bowel disease (Crohn's disease or ulcerative colitis), tell your healthcare provider if you have worsening disease symptoms during treatment with BIMZELX or develop new or worsening signs of Crohn's disease or ulcerative colitis.

### The most common side effects of BIMZELX for psoriatic arthritis include:

- upper respiratory tract infections
- headache
- urinary tract infections
- oral thrush or infections in the mouth
- diarrhea

These are not all of the possible side effects of BIMZELX. Call your doctor for medical advice about side effects.

Use BIMZELX exactly as your doctor tells you to use it.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.

Please see full Prescribing Information, including Patient Information, for BIMZELX available at BIMZELX.com.



Inspired by **patients.**  
Driven by **science.**

 **Bimzelx**<sup>®</sup>  
(bimekizumab-bkzx)